

Corrigendum to “Fixed effects and random effects estimation of higher-order spatial autoregressive models with spatial autoregressive and heteroskedastic disturbances.”

(WU Economics Working Paper, No 173, April 2014)

Harald Badinger and Peter Egger

September 2016

We are grateful to Di Liu for pointing out two errors in the definition of the matrices $\mathbf{A}_{1,N}^{s,s'}$ and $\mathbf{A}_{3,N}^{s,s'}$, appearing in the quadratic form of the moment conditions in equations (12a) and (12c).

In equation (12a), the correct definition of matrix $\mathbf{A}_{1,N}^{s,s'}$ is

$$\mathbf{A}_{1,N}^{s,s'} = \mathbf{Q}_{0,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N}) - \text{diag}_{n=1}^{NT} [\mathbf{Q}_{0,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N})_{nn}].$$

In equation (12c), the correct definition of matrix $\mathbf{A}_{3,N}^{s,s'}$ is

$$\mathbf{A}_{3,N}^{s,s'} = \mathbf{Q}_{1,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N}) - \text{diag}_{n=1}^{NT} [\mathbf{Q}_{1,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N})_{nn}].$$

The definition of the GM estimator (equations (13)-(18)) does not make use of the matrices $\mathbf{A}_{1,N}^{s,s'}$ and $\mathbf{A}_{3,N}^{s,s'}$ and remains unchanged. The only consequential error appears in the derivation of the variance-covariance matrix of the GM estimator, where the matrices $\mathbf{A}_{1,N}^{s,s'}$ and $\mathbf{A}_{3,N}^{s,s'}$ enter through equation (22) and show up ultimately in the definition of the blocks of the matrices $\bar{\mathbf{A}}_{1,N}^{s,s'}$ and $\bar{\mathbf{A}}_{3,N}^{s,s'}$ in equations (26a)-(26d).

In equation (26a), the correct definitions of the blocks of matrix $\bar{\mathbf{A}}_{1,N}^{s,s'}$ are

$$\begin{aligned} \mathbf{A}_{1,v,N}^{s,s'} &= \frac{1}{2(T-1)} [\mathbf{A}_{1,N}^{s,s'} + (\mathbf{A}_{1,N}^{s,s'})'] \\ \mathbf{A}_{1,\mu,N}^{s,s'} &= -\frac{1}{(T-1)} (\mathbf{e}'_T \otimes \mathbf{I}_N) \text{diag}_{n=1}^{NT} [\mathbf{Q}_{0,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N})] (\mathbf{e}_T \otimes \mathbf{I}_N) \\ \mathbf{A}_{1,v,\mu,N}^{s,s'} &= -\frac{1}{(T-1)} \text{diag}_{n=1}^{NT} [\mathbf{Q}_{0,N} (\mathbf{I}_T \otimes \mathbf{M}'_{s',N} \mathbf{M}_{s,N})] (\mathbf{e}_T \otimes \mathbf{I}_N). \end{aligned}$$

In equation (26c), the correct definitions of the blocks of matrix $\bar{\mathbf{A}}_{3,N}^{s,s'}$ are

$$\begin{aligned} \mathbf{A}_{3,v,N}^{s,s'} &= \frac{1}{2} \{ \mathbf{Q}_{1,N} [\mathbf{I}_T \otimes (\mathbf{M}'_{s',N} \mathbf{M}_{s,N} + \mathbf{M}'_{s,N} \mathbf{M}_{s',N})] \mathbf{Q}_{1,N} - 2 \text{diag}_{n=1}^{NT} [(\mathbf{Q}_{1,N} (\mathbf{I}_T \otimes (\mathbf{M}'_{s',N} \mathbf{M}_{s,N})))] \} \text{ or} \\ \mathbf{A}_{3,v,N}^{s,s'} &= \frac{1}{2} [\mathbf{A}_{3,N}^{s,s'} + (\mathbf{A}_{3,N}^{s,s'})'], \\ \mathbf{A}_{3,\mu,N}^{s,s'} &= \frac{1}{2} \{ T(\mathbf{M}'_{s',N} \mathbf{M}_{s,N} + \mathbf{M}'_{s,N} \mathbf{M}_{s',N}) - 2(\mathbf{e}'_T \otimes \mathbf{I}_N) \text{diag}_{n=1}^{NT} [(\mathbf{Q}_{1,N} (\mathbf{I}_T \otimes (\mathbf{M}'_{s',N} \mathbf{M}_{s,N})))] (\mathbf{e}_T \otimes \mathbf{I}_N) \}, \\ \mathbf{A}_{3,v,\mu,N}^{s,s'} &= \frac{1}{2} \{ [\mathbf{e}_T \otimes (\mathbf{M}'_{s',N} \mathbf{M}_{s,N} + \mathbf{M}'_{s,N} \mathbf{M}_{s',N})] - 2 \text{diag}_{n=1}^{NT} [(\mathbf{Q}_{1,N} (\mathbf{I}_T \otimes (\mathbf{M}'_{s',N} \mathbf{M}_{s,N})))] (\mathbf{e}_T \otimes \mathbf{I}_N) \}. \end{aligned}$$